



This booklet uses easy words and pictures.

You might want to ask someone to help you read it



You may not feel like having sex straight away after you have had your baby.

You may feel sore and bruised and it may feel uncomfortable to have sex.




As soon as you are ready to have sex again you can get pregnant.

You can also get or spread sexually transmitted infections.



It is important to think about contraception so you don't get pregnant when you don't want to.




Some contraception methods can protect you from getting or spreading sexually transmitted infections.



Breastfeeding can delay your periods but is not a contraceptive.



You can still get pregnant when breastfeeding.



You can still get pregnant even if you haven't had a period after your baby has been born.



Your midwife can arrange for you to have a contraceptive implant put in when you are in hospital after having your baby.

5



Talk to your midwife, health visitor or GP if you are worried about anything to do with contraception or sex after a baby.

This leaflet has been developed by Leeds Teaching Hospitals NHS Trust. It has kindly been shared for use by HNY Local Maternity and Neonatal System (LMNS).



© The Leeds Teaching Hospitals NHS Trust • 1st edition (Ver 1) Developed by: Laura Walton, Deputy Head of Midwifery All images / Resources Copyright © LYPFT easyonthei

Produced by: Medical Illustration Services • MID code: M20221205 013/BP

LN005508 Publication date 10/2023 Review date 10/2026